

NyGuard® IGR Concentrate is a juvenile hormone mimic which delivers strong IGR performance and allows you to effectively break the reproductive cycle of your toughest insects and related pests.

	Gentrol®	Precor®	NyGuard® IGR
Broadest spectrum (50+ pests)			✓
Labeled for mosquito control including Aedes that may carry and transmit Zika virus			✓
EPA reduced risk* status			✓
Food areas & non-food areas	✓		✓
Use indoors & outdoors			✓
Photostable			✓
Long-lasting residual activity	✓	✓	✓



Product Specifications

Signal Word	Caution
Packaging	140 mL bottles (10 per case), 480 mL bottles (6 per case)
EPA Registration Number	1021-1603
Food Handling	For use in food and non-food areas
Stability of Undiluted Product	Stable
Stability of Diluted Product	Do not allow to stand overnight. Mix before each use.
Appearance	Clear, colorless liquid
Odor	Sweet surfactant odor
Active Ingredients	Pyriproxyfen (Nylar)
Flammability	OSHA – combustible liquid
Mode of Action	Juvenile hormone mimic
Class of Chemistry	Insect growth regulator (IGR)
Respirator Required	Not required
Mix or Dilute in	Water or petroleum distillates
Activity	Residual – up to 7 months
Shelf Life	12 months closed in original commercial packaging

Use Areas

Indoor in Food and Non-Food Areas, such as:

- Homes
- Food processing plants
- Restaurants
- Schools
- Warehouses
- Hotels

Outdoor Perimeter Treatments to areas, such as:

- Doors
- Garage doors
- Patios
- Decks
- Carports and around garbage areas

In, On or Around Livestock and Poultry Housing, and Pet Kennels

*Pyriproxyfen, the active ingredient in NyGuard, has EPA Reduced Risk status in Food Handling.

Method of Application

- Broadcast surface treatment
- Crack and crevice or spot treatment
- ULV fogging
- Perimeter spray
- Power spray
- Tank mix or apply alone

Best Practices when using NyGuard® IGR Concentrate

Use NyGuard as part of an Integrated Pest Management (IPM) program to break the cycle of insect reproduction. NyGuard gets to the root cause of insect problems – the eggs, larvae or nymphs – and prevents them from maturing into reproductive adults. NyGuard will cause morphological abnormalities and disrupt the normal development of immature stages of insects, leading to sterility or death. When adults are present, use NyGuard in conjunction with a non-IGR insecticide. This method focuses on the large-scale removal of the reproductive adult component of the target population, while at the same time introducing NyGuard to disrupt the development of further generations.

Key Insects Controlled

Ants
Asian Cockroaches

Brown Banded Cockroaches
Carrion Beetles
Cigarette Beetles
Confused Flour Beetles
Crickets
Darkling Beetles
Dermestid Beetles
Fleas
Flies
Fungus Gnats
German Cockroaches
Hide Beetles
House Flies
Indian Meal Moths
Lesser Grain Borers
Litter Beetles
Merchant Grain Beetles
Midges
Mosquitoes
Moths
Red Flour Beetles
Rice Moths
Sawtooth Grain Beetles
Stable Flies
Tobacco Moths
Warehouse Beetles

How NyGuard® Breaks the Life Cycle of Your Toughest Insect Pests				
Complete metamorphosis Includes insects such as: Fleas, Moths, Beetles (including but not limited to Darkling Beetles, Carpet Beetles and Sawtoothed Grain Beetles), Ants, Flies, Midges, Mosquitoes and Fungus Gnats.				
	Egg stage Effects of NyGuard exposure: • Mortality	Larval stage Effects of NyGuard exposure: • Permanent larval stage • Reduced fitness • Increased susceptibility to non-IGR insecticides • Sterility in adult stage if development is able to continue	Pupal stage Results of NyGuard exposure in larvae stage: • Death • Morphological defects • No adult eclosion or emergence	Adult stage Results of NyGuard exposure: • Reduced egg production • Decreased viability of eggs
Incomplete metamorphosis Includes insects such as: Cockroaches and Crickets.				
	Egg stage Effects of NyGuard exposure: • Mortality	Nymphal stage Effects of NyGuard exposure: • Permanent nymphal stage • Reduced fitness • Increased susceptibility to non-IGR insecticides • Morphological defects • Behavioral changes—more active during daylight hours • Death • Sterility in adult stage if development is able to continue		Adult stage Results of NyGuard exposure: • Reduced reproduction—fewer eggs • Decreased viability of eggs

Always read and follow label and SDS directions.

To learn more, visit www.mgk.com, call 1-800-645-6466 or send an e-mail to brands@mgk.com.

